Ephestia cautella (Walk.)

(Pyralidae : Lepidoptera) Ephestia calidella (Guenee)

(2002/10/1 2002/7/24

Ephestia calidella (Guenee) Ephestia cautella (Walk.)

Match

% 650 100 Dimilin .Trigard

Effect of Host Kind and Egg Treatment by Sub-lethal **Concentration of Some Insect Growth Inhibitors on Some Biological** Characteristics of Fig Moth Ephestia cautella (Walk) and Currant Moth Ephestia calidella (Guenee) (Pyralidae : Lepidoptera)

Nazar M. Mallah

Renna R. Al-Sabie

Department of Plant Protection College of Agriculture and Forestry Mosul University

ABSTRACT

The results of this study revealed that the LC₅₀ values of Dimilin 10% E.C, Match 50% E.C and Trigard75% W.P. on eggs of Ephestia cautella (Walk.) and Ephestia

)

88

calidella (Guenee) were varied according to the host kinds and insect species. Match has a superior value of toxicity ratio and relative efficiency on eggs of *E. cautella* reaches 100 and 650 % respectively. The eggs of both species treated with sub-lethal concentration of insect growth inhibitors showed an increasing in eggs incubation period and reduction in eggs hatching percentage comprised with the control. The effect of egg treatment expand to the larval stage and increase the larval period, reduce the pupation percentage, elongate the pupal period, reduce the eclosion percentage of the adult and reduce the adult stage period and the number of eggs laid by the female.

Ephestia calidella (Guenee) Ephestia cautella (Walk.) .(1988 Ahmed, 1985) %19.86-7.58 (1987) (Attia, 1976 Hashimoto, 1964) (Schulten, 1970)) .(1987 2001 $%10 \pm 45$ °3 ±25

12

(

1

137

```
(Trigard 75% W.P Match 50% E.C Dimilin 10% E.C.)
                             % 1 0.8 0.5 0.3
                                   (
                                        48-24)
                                                                 20
                                       1
                                                   Potter Tower
                               .(Busvine, 1971) <sup>2</sup>
                                                      /
                                                             12
                                   (Abbott, 1925)
               LC_{50}
                                   .(Litchfield and Wilcoxon, 1949)
                                           .(Sun and Johnson, 1960)
                                                    LC50
                   100×
                                            LC50
                                                 LC50
               100×
                                         LC50
 )
                      0.003
                                                    (
                                                    (Trigard Match
                                                                       Dimilin)
                                       1
                                                   Potter Tower
                                            (Busvine, 1971) <sup>2</sup>
        30
                                                                        12
                                  9
                                                                  (48-24)
                                5
                                               3×6)
```

)) %5 .(1990 (1) (Trigard Match Dimilin) Match LC50 (0.0012)Trigard (1) (0.0078) LC50 Charmillot et al. (2001) .(0.0047) (0.0078-0.0012) C.pomonella(L.)Diflubenzuron Hexaflumuron Teflubenzuron 15 1.3 0.6

> %5 Match %100 Trigard %15.38

(1)

139

%5 (1)

%650

%100 Match

. Trigard

:1

%	%	-		LC50		
325	50	0.002-0.002	2.17	0.0024		Dimilin
210.80	32.43	0.007 -0.001	1.79	0.0037		Dillillilli
650	100	0.018-0.008	2.64	0.0012		Match
169.57	26.09	0.009-0.002	1.77	0.0046		Match
236.36	36.36	0.004-0.002	1.94	0.0033		Trigard
150	23.08	0.009-0.002	1.82	0.0052		Trigaru
173.30	26.67	0.004-0.004	2.41	0.0045		Dimilin
144.40	22.22	0.008-0.003	2.02	0.0054		Dillillilli
190.24	29.27	0.006-0.002	1.90	0.0041		Match
125.81	19.35	0.01-0.003	2.85	0.0062		Match
173.30	26.67	0.004-0.004	2.41	0.0045		Trigard
150	23.08	0.006-0.004	1.90	0.0052		Iligalu
114.71	17.65	0.014-0.003	1.87	0.0068		Dimilin
185.70	28.57	0.004-0.004	2.25	0.0042		Dillillill
144.40	22.22	0.013-0.002	1.64	0.0054		Match
162.50	25	0.008-0.002	1.84	0.0048		Match
100	15.38	0.003-0.001	2.74	0.0078		Trigard
139.26	21.42	0.009-0.003	1.95	0.0056		ingalu

.%5

140

: :

(2)

7. 5

Match 5.25
Dimilin Dimilin

Match

. 5 (2)

Match Trigard Dimilin 10 10.50 10.75

(2001)

(2) . Trigard

Carvalho (1994)

Triflumuron Buprofezin Trigard

(2)

141

:2

								•	
	1								
75	- 60	- 80	- 85	7.50	5.00	5.25	4.75	Dii	milin
75	- 80	- 75	- 70	7.50	10.50	10.75	8.75	Dil	
(1.66	50	- 75	- 65	7.29	5.25	5.00	4.75	N.A.	atch
61.66	55	- 60	- 65		- 10.00	- 10.00	8.75	IVI	atcn
0.46	- 60	- 80	- 70	6.62	4.50	4.25	4.50	Twi	aand
9.16	- 75	- 70	- 60		- 9.50	10.50	6.50	111	igard
91.66	- 85	90	95	6.50	4.50	4.25	4.25	Co	ntrol
91.00	95	95	90	0.50	9.25	- 9.75	7.00	Co	111101
	* 70	* 78.12	* 75		* 7.31	* 7.46	* 6.15		
			74.58	•		4.68			
			74.16		Γ	9.27			
			%	5		-	()*	
	. %5						()		

.

Match % 55 50 (2) (1998) Match 100 (1999)%69.4 Sesamia cretica (Led.) Match 200 (3) %5 55.5 (3) Match Dimilin 52 54.25 Match Trigard (1998) Match (2001) Trigard Trigard) ((3) Match Dimilin 7.45 40

%80

143

					/			
65.83	- 50	- 65	- 80	49.29	- 49.75	- 47.5	- 47.5	Dimili
05.65	- 60	- 60	- 80	49.29	- 49.5	54.25	- 47.75	Dillilli
52.5	40	- 55	- 60	51.45	- 50.0	- 49.25	- 50.5	Match
52.5	- 45	- 55	- 60		- 52.0	55.5	- 51.5	Mater
65	- 45	- 75	- 70	47.29	- 47.0	- 45.25	- 48.75	Trigar
05	- 65	- 65	- 70	41.29	- 47.25	- 52.0	- 43.5	Tilgar
87.5	- 80	90	90	37.70	33.5	38.25	30.5	Contro
07.5	85	90	90	37.70	40.5	- 49.25	34.25	Contro
	* 58.75	* 69.37	* 75		* 46.18	* 48.90	* 44.21	
			66.66	!		44.81	_	
			68.75		ſ	48.06		
		<u>.</u>		%5	_	<u> </u>	(()*

. %5

. %5

		15.5		(4)	
	13.75			Match	
			Match		
		. 15.5-8.25		%	65
		. 13.3 0.23	(4)		•
	Matah	(1997)			
	Match			(200	1)
(4)			5	7.2	Trigard
			·/ -		
			% 5		Makala
Trigard				%40 30	Match
IIIguiu	%45		% 40	7010 30	
				Dimilin	%70
%80-30					
,		(1227)			
)		(1997)	(
		%30.20	(97.6		
Pectinophora		Diflubenzuron		Flint et al. (1987 go	() ssypiella

145

.

45 - 55 - 60 9.75 9.75 11 _ 60.83 11.12 Dimilin - 65 - 10.75 - 13 70 - 70 - 13 30 - 12 - 11.5 - 50 45 - 12.75 45.83 13.08 Match 40 - 50 - 13 15.5 13.75 - 60 40 65 70 9.75 - 8.25 10.5 61.66 10.41 Trigard 45 - 70 80 - 11.5 10.5 - 12 80 85 6.75 6.75 85 85.83 8.45 Control 85 90 90 9.50 - 10 - 10.75 55 68.75 10.37 11.12 66.87 10.81 9.60 59.16 11.93 67.91 %5

%5

:4

(5) %50 3.75 3.5 Match Dimilin Trigard %5 (1997) Match Match (1998) (5) / %5 / 89 85 Dimilin Match 90 (1997) Match 98.5 128 (1988) Hofmeister

147

:5

1					·				
96.08	- 99.5	- 97.75	91.5	5.12	- 5.25	- 5.75	- 4.5		Dimilin
90.00	93.5	- 104.25	90	5.12	- 5.25	- 5.25	- 4.75		
95.16	- 98.75	- 98.75	- 10.2	4.2	- 4.25	3.50	3.75		Match
95.10	85	- 97.5	89	4.2	- 5	- 4.25	- 4.5		
105.58	- 108.25	- 180.25	- 97.5	F 25	- 5.25	- 4.75	- 4.75		Trigard
105.58	- 107.5	- 112	- 100	5.25	- 5.75	- 5.25	- 5.75		Trigard
130.66	137.5	- 122.25	152.25	6.66	- 6	- 6	- 6.5		Control
130.00	- 110.25	- 130.25	- 131.5	0.00	7.5	7	7		Control
	* 105.03	* 108.87	* 106.71		* 5.53	* 5.21	* 5.18		
	_		109.52			5.02			
		Ī	104.22			5.6			
				%5	-	•	()*	-

%5

.1998

Ephestia cautella (Walk..)

Lufenuron (Match)

(Pyralidae:Lepidoptera)

.85-81 2

Musca .2001

.(Muscidae : Diptera) domestica (L.)

.1990

Ephestia cautella (Walk..) .1987

.

. Match .1997

•

Lepidoptera:) Sesamia cretica (Led.)

.1999

1 17 . (Phalaenidae

.8-1

. Alsystin .1988

Abbott, WS., 1925. A method for computing the effectiveness of an insecticide. J. Econ. Ent., Vol.18, pp.265-267.

Ahmed, M.S.H., S.K. Al-Maliky, A.A. Al-Tawoel, N.F. Jabo, Z.S., Al-Hakkak, 1985. Effect of three temperatures regimes on rearing and biological activaties of *Bracon hebator*. J. Stored Prod. Res., Vol.21, Np.2, pp.65-68.

Attia, F.I., 1976. Insecticide resistance in *Cadra cautella* in new South Wales, Australia. J. Econ. Ent., Vol.66, pp.773–774.

Busvine, J.R., 1971. A critical review of the techniques for testing insecticides, 2nd ed. Commonwealth Agricultural Bureaux, 345p.

Carvalho, G.A. 1994. Effects of insect growth regulators on eggs and larvae of *Ceraeochrysa cubana*. Drazal. Cienc. Prat., Vol.18, No.1, pp.49-55.

- Charmillot, P.J., A. Gourmelom, A.L. Fabre, D. Pasquier, 2001. Ovicidal and larvicidal effectiveness of several growth inhibitors and regulators on the codling moth *Cydia pomonella* (L.) (Lepidoptera: Tortricidae). J. Appl. Ent., Vol.125, No.3, pp.147-153.
- Flint, H.M., R.L. Smith, J.M. Noble, D. Shaw, A.B. Demilo and Khalil, 1987. Laboratory tests of diflubenzuron and four analogues against the pink bollworn and afield cage test with diflubenzuron and EI-494 for control of the pink bollworn and cotton leaf hopper. J. Econ. Ent., Vol.71, pp.616-619.
- Hashimoto, Y., 1964. Resistance to insecticides of almond moth, *Ephestia cautella*. I. Development of methylparathion resistince. Konchu-Gaku Zasshi, Vol.8, No.1, pp.62-68.
- Hofmeister, P.C. Kunast, A. Lang, 1988. N. Benzoyl-N-Phenoxyphenyl and N-Benzoyl-N-Carboxyphenylureas: A review of their chemical synthesis and boilogical profites. Pestic. Sci., Vol.22, pp.221-230.
- Litchfield, J.R. and F. Wilcoxon, 1949. A simplified method of evaluating dose effect experiments. J. Pharmacology and Experimental Therapy, Vol.96, pp.99-113.
- Schulten, G.M., 1970. Preliminary results of field trails on control of *Ephestia cautella* as apest of bagged maize in malawi, PANS-, Vol.16, No.4, pp.709-713.
- Sun, Y.P. and E.R. Johnson, 1960. Synergistic and antagonistic actions of insecticide-synergist combinations and their mode of action. J. Agric. Food. Chem., Vol.8, No.4, pp.261-266.